

SPLITTING IN THE K -THEORY LOCALIZATION SEQUENCE OF NUMBER FIELDS

Luca Caputo*

February 25, 2010

ABSTRACT - Let p be a rational prime and let F be a number field. Then, for each $i \geq 1$, there is an exact localization sequence

$$0 \longrightarrow K_{2i}(\mathcal{O}_F)_p \longrightarrow K_{2i}(F)_p \xrightarrow{\partial_{F,i}} \bigoplus_{v \nmid \infty} K_{2i-1}(k_v)_p \longrightarrow 0$$

where $K_j(R)$ denotes the j -th Quillen's K -theory group of the ring R . If p is odd or F is nonexceptional, we find necessary and sufficient conditions for this exact sequence to split: these conditions involve coinvariants of twisted p -parts of the p -class groups of certain subfields of the fields $F(\mu_{p^n})$ for $n \in \mathbb{N}$. We also compare our conditions with the weaker condition $WK_{2i}^{\acute{e}t}(F) = 0$ and give some example.

1 Introduction and notation

Throughout the paper, p will denote a rational prime. For an abelian group A , set

$$\text{Div}(A) = \text{maximal divisible subgroup of } A$$

$$\text{div}(A) = \{a \in A \mid \forall n \in \mathbb{N} \exists a_n \in A : a = na_n\}$$

Then $\text{div}(A)$ is a subgroup of A which is commonly called the subgroup of (infinitely) divisible elements or the subgroup of elements of infinite height of A . We denote by A_p the p -primary part of A and, for $n \in \mathbb{N}$, by $A[p^n]$ the subgroup of elements of A whose order divides p^n .

If R is a ring and j is a natural number, $K_j(R)$ denotes the j -th Quillen's K -group of R . Let F be a number field and i be a positive integer. Thanks to Soulé's results (see [21], Theorem 4.6), Quillen's long exact localization sequence splits into isomorphisms $K_{2i+1}(\mathcal{O}_F) \cong K_{2i+1}(F)$ and short exact sequences of the form

$$0 \longrightarrow K_{2i}(\mathcal{O}_F) \longrightarrow K_{2i}(F) \xrightarrow{\partial_{F,i}} \bigoplus_{v \nmid \infty} K_{2i-1}(k_v) \longrightarrow 0 \quad (1)$$

where $\mathcal{O}_F$ is the ring of integers of F , k_v is the residue field of F at v and the direct sum is taken over the finite places of F . We recall that, thanks to Quillen's and Borel's results, $K_{2i-1}(k_v)$ is cyclic of order $|k_v|^i - 1$ and $K_{2i}(\mathcal{O}_F)$ is a finite group: in particular we always have $\text{Div}(K_{2i}(F)) = 0$ (but in general $\text{div}(K_{2i}(F))$ may be nontrivial).

One can ask for conditions for the exact sequence in (1) to split. As a motivation for this question we quote the following three results:

- if $F = \mathbb{Q}$, Tate showed that the exact sequence (1) splits (see [11], Theorem 11.6);
- if E is a rational function field in one variable over an arbitrary base field, then the localization sequence for $K_2(E)$ (which is completely analogous to (1)) always splits, thanks to a result of Milnor and Tate (see [10], Theorem 2.3);

*Partially supported by an IRCSET fellowship.

- if E is a local field (i.e. a field complete with respect to a discrete valuation whose residue field is finite), then the localization sequence for $K_{2i}(E)$ (which again is completely analogous to (1)) always splits, thanks to a result of Soulé (see [17], Proposition 4).

Coming back to the case of a number field F , consider now the exact sequence which (1) induces on p -primary parts, which we shall refer to as the localization sequence for $K_{2i}(F)_p$. The problem of the splitting of the localization sequence for $K_{2i}(F)_p$ was first studied by Banaszak: in one of his papers (see [1], Corollary 1) he claims that, if p is odd, the localization sequence for $K_{2i}(F)_p$ splits if and only if $\text{div}(K_{2i}(F))_p = 0$ (in fact Banaszak's result is stated in terms of $WK_{2i}^{\text{ét}}(F)$, see Proposition 2). This is obviously a necessary condition, since both the right and the left terms of the localization sequence for $K_{2i}(F)_p$ have trivial subgroup of divisible elements. However the proof of sufficiency seems to be incomplete. It turns out that, for any $i \geq 1$, there is a counterexample, namely there is a number field F and a prime p such that $\text{div}(K_{2i}(F))_p = 0$ but the localization sequence for $K_{2i}(F)_p$ does not split (see Example 2). The counterexample is constructed using Theorem 1, which is our main result and is described in Section 3: under the assumption that p is odd or F is nonexceptional (see Definition 1), the obstruction to the existence of a splitting for the localization sequence for $K_{2i}(F)_p$ are the p -parts of the p -class groups $Cl_{F_{i,n}}^{S_{i,n}}$ of certain subfields $F_{i,n}$ of the fields $F(\mu_{p^n})$, where μ_{p^n} denotes the group of roots of unity in an algebraic closure of F . The main idea of the proof of Theorem 1 is essentially to use a generalization of an exact sequence of Keune which builds on an earlier result of Tate (Proposition 3). For instance, if $\mu_{p^n} \subseteq F$, a consequence of this exact sequence (which has already been used in [4]) is an isomorphism

$$(Cl_F^S)_p \otimes \mu_{p^n} \cong (K_2(\mathcal{O}_F) \cap p^n K_2(F)) / p^n K_2(\mathcal{O}_F)$$

and the triviality of the left-hand term is closely related to the splitting of the localization sequence for $K_2(F)_p$. Using a codescent argument, the hypothesis $\mu_{p^n} \subseteq F$ can be removed and this can be generalized to $K_{2i}(F)$ but one has to face the absence of a description of $K_{2i}(F)$ analogous to that of Matsumoto for $K_2(F)$. The way we found to overcome this difficulty is to work with continuous Galois cohomology instead of K -theory (the connection between the two theories is classical). This translation implies no further assumptions as explained in Section 2, which is essentially taken from [1], with some slight change and generalization.

The difference between our splitting criterion and the condition $\text{div}(K_{2i}(F))_p = 0$ (which is equivalent to the vanishing of the i -th étale wild kernel $WK_{2i}^{\text{ét}}(F)$) is also analyzed at the end of Section 3. Anyway the condition $\text{div}(K_{2i}(F))_p = 0$ is often necessary and sufficient for the localization sequence for $K_{2i}(F)_p$ to split (for example in the case where $F = \mathbb{Q}$, see Example 1).

Acknowledgements

I wish to thank Kevin Hutchinson, Jean-François Jaulent and Manfred Kolster for their suggestions and comments.

2 Localization sequence for continuous Galois cohomology

In this section we are going to translate the problem of the existence of a splitting for the localization sequence for $K_{2i}(F)_p$ in cohomological terms. First of all we recall the following notion.

Definition 1. Let E be a field of characteristic other than 2. Then E is said to be *nonexceptional* if $\text{Gal}(E(\mu_{2^\infty})/E)$ has no element of order 2 (and *exceptional* otherwise). Here μ_{2^∞} is the group of roots of unity whose order is a power of 2 in an algebraic closure of E .

Remark 1. Note that nonexceptional fields have no embeddings in $\mathbb{R}$ (since $\mathbb{R}$ is exceptional and subfields of exceptional fields are exceptional).

As in the preceding section, p denotes a rational prime, i is a positive integer and F is a number field.

We are interested in the localization sequence for $K_{2i}(F)_p$, namely

$$0 \longrightarrow K_{2i}(\mathcal{O}_F)_p \longrightarrow K_{2i}(F)_p \xrightarrow{\partial_{F,i}} \bigoplus_{v \nmid p\infty} K_{2i-1}(k_v)_p \longrightarrow 0 \quad (2)$$

which easily follows from (1). This exact sequence has (at least conjecturally, see Remark 2) a cohomological counterpart, as shown in the next proposition (which can essentially be found [1], Section I, §2-3). In this paper, cohomology is always continuous cohomology of profinite groups, in the sense of Tate ([19]).

Notation 1. If T is a finite set of places of F , $G_{F,T}$ denotes the Galois group of the maximal extension F_T of F unramified outside T and $\mathcal{O}_F^T$ is the ring of T -integers of F . For a field E , G_E denotes the Galois group of a separable algebraic closure of E (and we use the convention $H^j(E, -) := H^j(G_E, -)$ for cohomology). Finally, $S_F^{(p)} = S$ denotes the set of primes above p and ∞ in F .

For a noetherian $\mathbb{Z}[\frac{1}{p}]$ -algebra A and a natural number j , let $K_j^{\text{ét}}(A)$ denote the j -th étale K -theory group of Dwyer and Friedlander (see [3]). For any finite set T containing S , there are functorial maps

$$K_{2i}(\mathcal{O}_F^T)_p \xrightarrow{\nu} K_{2i}^{\text{ét}}(\mathcal{O}_F^T) \quad \text{and} \quad K_{2i}^{\text{ét}}(\mathcal{O}_F^T) \xrightarrow{\alpha} H^2(G_{F,T}, \mathbb{Z}_p(i+1))$$

We know that α is an isomorphism (see [3], Remark 8.8). We set $ch = \alpha \circ \nu$ (ch stands for Chern character).

Proposition 1 (Banaszak). *Suppose that p is odd or F is nonexceptional. There is a commutative diagram with exact rows as follows*

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_{2i}(\mathcal{O}_F^S)_p & \longrightarrow & K_{2i}(F)_p & \xrightarrow{\partial_{F,i}} & \bigoplus_{v \nmid p\infty} K_{2i-1}(k_v)_p \longrightarrow 0 \\ & & \downarrow ch & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H^2(G_{F,S}, \mathbb{Z}_p(i+1)) & \longrightarrow & H^2(F, \mathbb{Z}_p(i+1))_p & \xrightarrow{\partial_{F,i}^c} & \bigoplus_{v \nmid p\infty} H^1(k_v, \mathbb{Z}_p(i)) \longrightarrow 0 \end{array}$$

Moreover vertical maps in the above diagram are surjective and the rightmost is an isomorphism.

Proof. Let T be any finite set of primes T containing S . For the rest of this proof, for any $j \in \mathbb{N}$, set

$$H_T^j(-) := H^j(G_{F,T}, -) \quad \text{and} \quad H^j(-) := H^j(F, -)$$

Then, for any $n \in \mathbb{N}$, there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_{2i}(\mathcal{O}_F^S)_p & \longrightarrow & K_{2i}(\mathcal{O}_F^T)_p & \xrightarrow{\partial_{F,i}} & \bigoplus_{v \in T \setminus S} K_{2i-1}(k_v)_p \longrightarrow 0 \\ & & \downarrow ch & & \downarrow ch & & \downarrow \\ 0 & \longrightarrow & H_S^2(\mathbb{Z}_p(i+1)) & \longrightarrow & H_T^2(\mathbb{Z}_p(i+1)) & \xrightarrow{\partial_{F,i}^c} & \bigoplus_{v \in T \setminus S} H^1(k_v, \mathbb{Z}_p(i)) \longrightarrow 0 \\ & & \uparrow \delta & & \uparrow \delta & & \uparrow \delta \\ 0 & \longrightarrow & H_S^1(\mathbb{Q}_p/\mathbb{Z}_p(i+1))/\text{Div} & \longrightarrow & H_T^1(\mathbb{Q}_p/\mathbb{Z}_p(i+1))/\text{Div} & \longrightarrow & \bigoplus_{v \in T \setminus S} H^0(k_v, \mathbb{Q}_p/\mathbb{Z}_p(i)) \longrightarrow 0 \end{array}$$

The definition and the exactness of the middle row can be found in [17], Section III and and [15], Section 4. The upper rightmost vertical map is defined by the diagram (note that in general this does *not* coincide with the direct sum of the residue fields Chern characters) and it is bijective. This follows from the surjectivity of the ch 's (see [3], Theorem 8.7, for the case p odd and [15], Theorem 0.1, for the case $p = 2$ and F nonexceptional) together with the easy fact that, for any $v \in T \setminus S$, $H^1(k_v, \mathbb{Z}_p(i))$ is cyclic of order $|k_v|^i - 1$ and has therefore the same order as $K_{2i-1}(k_v)_p$ (thanks to Quillen's calculation). As for

the maps denoted with δ , they are connecting homomorphisms in the long exact cohomology sequence relative to the exact sequence

$$0 \rightarrow \mathbb{Z}_p(j) \rightarrow \mathbb{Q}_p(j) \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(j) \rightarrow 0$$

($j = i, i + 1$) and they are bijective (see [19], Proposition 2.3).

Taking direct limits as $T \supset S$ grows, we get

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_{2i}(\mathcal{O}_F^S)_p & \longrightarrow & K_{2i}(F)_p & \xrightarrow{\partial_{F,i}} & \oplus_{v \nmid p\infty} K_{2i-1}(k_v)_p \longrightarrow 0 \\ & & \downarrow ch\delta^{-1} & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H_S^1(\mathbb{Q}_p/\mathbb{Z}_p(i+1))/\text{Div} & \longrightarrow & H^1(\mathbb{Q}_p/\mathbb{Z}_p(i+1))/\text{Div} & \xrightarrow{\partial_{F,i}^c} & \oplus_{v \nmid p\infty} H^0(k_v, \mathbb{Q}_p/\mathbb{Z}_p(i)) \longrightarrow 0 \end{array} \quad (3)$$

This is because, for any finite set T containing S ,

$$\text{Div}(H_T^1(\mathbb{Q}_p/\mathbb{Z}_p(i+1))) = \text{Div}(H^1(\mathbb{Q}_p/\mathbb{Z}_p(i+1)))$$

(see [16], §4, Lemma 5). Composing again with δ we get a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_{2i}(\mathcal{O}_F^S)_p & \longrightarrow & K_{2i}(F)_p & \xrightarrow{\partial_{F,i}} & \oplus_{v \nmid p\infty} K_{2i-1}(k_v)_p \longrightarrow 0 \\ & & \downarrow ch & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H_S^2(\mathbb{Z}_p(i+1)) & \longrightarrow & H^2(\mathbb{Z}_p(i+1))_p & \xrightarrow{\partial_{F,i}^c} & \oplus_{v \nmid p\infty} H^1(k_v, \mathbb{Z}_p(i)) \longrightarrow 0 \end{array}$$

Here again the rightmost map is an isomorphism and the other two vertical maps are surjective. $\square$

Remark 2. The Quillen-Lichtenbaum conjecture predicts that for any finite set of places T containing S ,

$$ch : K_{2i}(\mathcal{O}_F^T)_p \longrightarrow H^2(G_{F,T}, \mathbb{Z}_p(i+1))$$

is indeed an isomorphism. Tate (see [19], Theorem 5.4) proved this statement for $i = 1$: in this case ch is just the Galois symbol ([3], proof of Theorem 8.2). The general case of the Quillen-Lichtenbaum conjecture is a consequence of the Bloch-Kato conjecture on the Galois symbol whose proof should appear soon.

In the following we will be mainly interested in the bottom row of the diagram of Proposition 1, namely the exact sequence

$$0 \rightarrow H^2(G_{F,S}, \mathbb{Z}_p(i+1)) \rightarrow H^2(F, \mathbb{Z}_p(i+1))_p \rightarrow \bigoplus_{v \nmid p\infty} H^1(k_v, \mathbb{Z}_p(i)) \rightarrow 0 \quad (4)$$

We will refer to it as the localization sequence for $H^2(F, \mathbb{Z}_p(i+1))$. The next proposition shows that, even without using the Quillen-Lichtenbaum conjecture, for our purposes there is no difference between considering (2) or (4).

Definition 2. A subgroup B of an abelian group A is *pure* if $nA \cap B = nB$ for each $n \in \mathbb{N}$.

Proposition 2. *Suppose that p is odd or F is nonexceptional. Then the localization sequence for $H^2(F, \mathbb{Z}_p(i+1))$ splits if and only if the localization sequence for $K_{2i}(F)_p$ splits.*

Proof. A diagram chasing in the diagram of Proposition 1 shows that if the localization sequence for $K_{2i}(F)_p$ splits then the localization sequence for $H^2(F, \mathbb{Z}_p(i+1))$ splits. As for the converse, we can assume $p \neq 2$ because if $p = 2$, then F is totally imaginary and the ch 's are isomorphisms by [15], Theorem 0.1¹. Banaszak (see [1], Proposition 2) proved that the natural map

$$\nu : K_{2i}(\mathcal{O}_F)_p \longrightarrow K_{2i}^{\text{ét}}(\mathcal{O}_F[\frac{1}{p}]) \quad (5)$$

¹It is likely that there exists a proof of the proposition for $p = 2$ and F nonexceptional which does not use the results in [15].

is split surjective. We are going to prove the analogous result for the map

$$\nu : K_{2i}(F)_p \longrightarrow K_{2i}^{\acute{e}t}(F)_p \quad (6)$$

with the same strategy as Banaszak, taking into account that the groups involved are no longer finite (but still torsion). First of all, for each $n \in \mathbb{N}$, there is a commutative diagram

$$\begin{array}{ccccc} K_{2i+1}(F, \mathbb{Z}/p^n\mathbb{Z}) & \longrightarrow & K_{2i}(F)[p^n] & \longrightarrow & 0 \\ \downarrow & & \downarrow \nu & & \\ K_{2i+1}^{\acute{e}t}(F, \mathbb{Z}/p^n\mathbb{Z}) & \longrightarrow & K_{2i}^{\acute{e}t}(F)[p^n] & \longrightarrow & 0 \end{array}$$

with exact rows and surjective vertical maps (see [1], Diagram 1.6). This implies that the kernel C_i of the map ν in (6) is a pure subgroup of $K_{2i}(F)_p$. Moreover C_i is finite since it is easily seen to coincide with the kernel of the map in (5) (see [1], Remark 7). Hence Theorem 7 of [7] tells us that the map in (6) is split. By definition of the maps ch , this proves that the map

$$K_{2i}(F)_p \longrightarrow H^2(F, \mathbb{Z}_p(i+1))_p$$

in the diagram of Proposition 1 splits. The proof of the proposition is then achieved by a diagram chasing in the diagram of Proposition 1. $\square$

3 Main result

We are going to describe the obstruction to the existence of a splitting for the localization sequence for $H^2(F, \mathbb{Z}_p(i+1))$ in terms of coinvariants of twisted p -parts of the class groups of certain subfields of the fields $F(\mu_{p^n})$. For a field E , we denote by $\mu_{p^n}(E)$ the group of p^n -th roots of unity in an algebraic closure of E (the reference to E in $\mu_{p^n}(E)$ will be often omitted).

Notation 2. For typographical convenience, we set

$$\Omega_{F,i,n}^{(p)} = \Omega_{i,n}^{(p)} = (H^2(G_{F,S}, \mathbb{Z}_p(i+1)) \cap p^n H^2(F, \mathbb{Z}_p(i+1))_p) / p^n H^2(G_{F,S}, \mathbb{Z}_p(i+1))$$

Remark 3. Note that, from the definition of $\Omega_{i,n}^{(p)}$, we have $\Omega_{i,0}^{(p)} = 0$.

The following lemma shows that the $\Omega_{i,n}^{(p)}$'s are the obstructions to the existence of a splitting for the localization sequence for $H^2(F, \mathbb{Z}_p(i+1))$.

Lemma 1. *The localization sequence for $H^2(F, \mathbb{Z}_p(i+1))$ splits if and only if for every $n \in \mathbb{N}$ we have $\Omega_{i,n}^{(p)} = 0$.*

Proof. Thanks to Theorem 5 in [7] and the fact that any direct summand of an abelian group is pure, we have the following equivalences

$$\begin{aligned} \Omega_{i,n}^{(p)} = 0 \quad \forall n \in \mathbb{N} & \Leftrightarrow H^2(G_{F,S}, \mathbb{Z}_p(i+1)) \text{ is pure in } H^2(F, \mathbb{Z}_p(i+1))_p \\ & \Leftrightarrow \text{the localization sequence for } H^2(F, \mathbb{Z}_p(i+1)) \text{ splits.} \end{aligned}$$

$\square$

We will make use of the following notation (see [22]).

Notation 3. Let E be any field. If M is a G_E -module, we denote by $E(M)$ the fixed field of the kernel of the homomorphism $G_E \rightarrow \text{Aut}(M)$ induced by the action of $\text{Gal}(\overline{E}/E)$ on M .

For each $n \in \mathbb{N}$, we now introduce the subfield $F_{i,n}$ of $F(\mu_{p^n})$ which will be relevant for us. Such subfields has been used for the first time by Weibel in [22].

Notation 4. Let F be a number field. For $n \in \mathbb{N}$ and $i \geq 1$, set $F_{i,n} = F(\mu_{p^n}(F)^{\otimes i})$ (note that $F_{i,0} = F$ for any $i \geq 1$). We will also use the notation $\Gamma_{i,n} = \text{Gal}(F_{i,n}/F)$. If w is a place in $F_{i,n}$, then denote by $(k_{i,n})_w$ the residue field of $F_{i,n}$ at w (thus $(k_{i,0})_w = k_w$). Finally, let $S_{i,n}$ be the set of primes of $F_{i,n}$ above p and ∞ (thus $S_{i,0} = S$) and let $G_{S_{i,n}}$ denote the Galois group of the maximal extension of $F_{i,n}$ unramified outside $S_{i,n}$ (thus $G_{S_{i,0}} = G_S = G_{F,S}$). Of course, in all this notation, a reference to p should appear but the context should prevent any misunderstanding.

The following lemmas are technical ingredients of the proof of Theorem 1. Most of them are well-known, but we collect them for the convenience of the reader, giving proofs whenever we could not find an adequate reference in the literature.

Lemma 2. *Suppose that p is an odd prime or F is nonexceptional. Let $v \nmid p$ be a place in F . For every $n, m \in \mathbb{N}$, there are isomorphisms of $\Gamma_{i,m}$ -modules*

$$\bigoplus_{w|v} H^1((k_{i,m})_w, \mathbb{Z}_p(i))[p^n] \cong \bigoplus_{w|v} H^0((k_{i,m})_w, \mu_{p^n}^{\otimes i})$$

and

$$H^2(G_{S_{i,m}}, \mathbb{Z}_p(i))/p^n \cong H^2(G_{S_{i,m}}, \mu_{p^n}^{\otimes i})$$

Proof. Both assertions come from the cohomology sequence corresponding to the exact sequence

$$0 \rightarrow \mathbb{Z}_p(i) \xrightarrow{p^n} \mathbb{Z}_p(i) \rightarrow \mu_{p^n}^{\otimes i} \rightarrow 0$$

In fact

- (i) $H^0((k_{i,m})_w, \mathbb{Z}_p(i)) = 0$ for any place $w|v$ in F_m (since $i > 0$);
- (ii) $H^3(G_{S_{i,m}}, \mathbb{Z}_p(i)) = 0$ since $G_{S_{i,m}}$ has p -cohomological dimension less or equal to 2 (see [12], (8.3.17) and (8.3.18)): this is true even in the case $p = 2$ because in that case $F_{i,m}$ has to be nonexceptional and therefore it has no real embeddings.

Now, since conjugation in cohomology commutes with the connecting homomorphism, we have that, for any fixed $w_0|v$ in $F_{i,m}$, (i) gives an isomorphism of D_v -modules (D_v being the decomposition group at v in $F_{i,m}/F$)

$$H^1((k_{i,m})_{w_0}, \mathbb{Z}_p(i))[p^n] \cong H^0((k_{i,m})_{w_0}, \mu_{p^n}^{\otimes i})$$

and (ii) gives an isomorphism of $\Gamma_{i,m}$ -modules

$$H^2(G_{S_{i,m}}, \mathbb{Z}_p(i))/p^n \cong H^2(G_{S_{i,m}}, \mu_{p^n}^{\otimes i})$$

Furthermore

$$\bigoplus_{w|v} H^1((k_{i,m})_w, \mathbb{Z}_p(i))[p^n] = \text{Ind}_{D_v}^{\Gamma_m} H^1((k_{i,m})_{w_0}, \mathbb{Z}_p(i))[p^n]$$

for any fixed $w_0|v$ in $F_{i,m}$ and

$$\bigoplus_{w|v} H^0((k_{i,m})_w, \mu_{p^n}^{\otimes i}) = \text{Ind}_{D_v}^{\Gamma_n} H^0((k_{i,m})_{w_0}, \mu_{p^n}^{\otimes i})$$

This concludes the proof. □

Lemma 3. *Let $v \nmid p$ be a finite place of F . If w is a place of $F_{i,n}$ above v , then $(k_{i,n})_w = k_v(\mu_{p^n}^{\otimes i})$.*

Proof. Both $(k_{i,n})_w$ and $k_v(\mu_{p^n}^{\otimes i})$ are subextensions of $k_v(\mu_{p^n})/k_v$. Set D_v (resp. D'_v) for the decomposition group of v in $F_{i,n}/F$ (resp. $F(\mu_{p^n})/F$) and P for the Galois group of $F(\mu_{p^n})/F_{i,n}$. Identifying $\text{Gal}(k_v(\mu_{p^n})/k_v)$ with D'_v we get

$$\text{Gal}(k_v(\mu_{p^n})/k_v(\mu_{p^n}^{\otimes i})) = D'_v \cap P$$

$$\text{Gal}((k_{i,n})_w/k_v) = D_v = D'_v/D'_v \cap P$$

which proves the result. $\square$

Lemma 4. *Let G be a finite cyclic group and let M be a faithful G -module which is cyclic of order a power of p as an abelian group. Then*

- if $p \neq 2$, then $H_j(G, M) = 0$ for any $j \geq 1$ (or equivalently the norm map $M_G \rightarrow M^G$ is surjective);
- if $p = 2$ and $G \neq \{\pm 1\}$ (i.e. G is not a group of order 2 whose generator g satisfies $gm = -m$ for any $m \in M$), then $H_j(G, M) = 0$ for any $j \geq 1$ (or equivalently the norm map $M_G \rightarrow M^G$ is surjective);

Proof. See [22], Lemma 3.2 and Remark 3.2.1. $\square$

Lemma 5. *Let $v \nmid p$ be a finite place of F . If w is a place of $F_{i,n}$ above v , then the corestriction homomorphisms*

$$H^0((k_{i,n})_w, \mu_{p^n}^{\otimes i}) \rightarrow H^0(k_v, \mu_{p^n}^{\otimes i})$$

are surjective if p is odd or n is large enough.

Proof. We have

$$H^0((k_{i,n})_w, \mu_{p^n}^{\otimes i}) = \mu_{p^n}^{\otimes i} \quad \text{and} \quad H^0(k_v, \mu_{p^n}^{\otimes i}) = H^0((k_{i,n})_w, \mu_{p^n}^{\otimes i})^{\text{Gal}((k_{i,n})_w/k_v)} = (\mu_{p^n}^{\otimes i})^{\text{Gal}((k_{i,n})_w/k_v)}$$

for any $n \in \mathbb{N}$. In particular the corestriction coincides with the norm $\mu_{p^n}^{\otimes i} \rightarrow (\mu_{p^n}^{\otimes i})^{\text{Gal}((k_{i,n})_w/k_v)}$. We then conclude by Lemma 4 with $M = \mu_{p^n}^{\otimes i}$ and $G = \text{Gal}((k_{i,n})_w/k_v)$ whose hypotheses are satisfied since $\text{Gal}((k_{i,n})_w/k_v)$ is always cyclic (since finite fields of odd characteristic are nonexceptional), acts faithfully on $\mu_{p^n}^{\otimes i}$ by Lemma 3 and is different from $\{\pm 1\}$ if n is large enough. $\square$

Lemma 6. *Let $c : \bigoplus_{w|p} \mu_{p^n}^{\otimes i}(F) \rightarrow \mu_{p^n}^{\otimes i}(F)$ be the codiagonal map $(\zeta_w)_w \mapsto \prod_w \zeta_w$. Then c is a surjective map of $\Gamma_{i,n}$ -modules and, for any $j \geq 1$, $H_j(\Gamma_{i,n}, \text{Ker } c) = 0$ if p is odd or n is large enough.*

Proof. If D_v denotes the decomposition group at v in $F_{i,n}/F$, we have

$$H_j(\Gamma_{i,n}, \bigoplus_{w|p} \mu_{p^n}^{\otimes i}) = H_j(\Gamma_{i,n}, \bigoplus_{v|p} \text{Ind}_{D_v}^{\Gamma_{i,n}} \mu_{p^n}^{\otimes i}) = \bigoplus_{v|p} H_j(D_v, \mu_{p^n}^{\otimes i})$$

by Shapiro's lemma, for any $j \geq 0$. Moreover if p is odd or n is large enough

$$H_j(\Gamma_{i,n}, \mu_{p^n}^{\otimes i}) = 0 = H_j(D_v, \mu_{p^n}^{\otimes i})$$

for any $j \geq 1$ by Lemma 4 and the result follows by considering the $\Gamma_{i,n}$ -homology sequence of the exact sequence

$$0 \rightarrow \text{Ker } c \rightarrow \bigoplus_{w|p} \mu_{p^n}^{\otimes i}(F) \xrightarrow{c} \mu_{p^n}^{\otimes i}(F) \rightarrow 0$$

$\square$

Lemma 7. *Suppose that p is an odd prime or F is nonexceptional. For every $n \in \mathbb{N}$, the corestriction map induces an isomorphism*

$$H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes i})_{\Gamma_{i,n}} \xrightarrow{\sim} H^2(G_S, \mu_{p^n}^{\otimes i})$$

Proof. Apply [22], Proposition 2.2, using the fact that G_S has p -cohomological dimension less or equal to 2 (see the proof of Lemma 2). $\square$

Notation 5. For any $i \geq 1$ and any $n \in \mathbb{N}$, we will denote by $Cl_{F_{i,n}}^{S_{i,n}}$ the class group of $\mathcal{O}_{F_{i,n}}^{S_{i,n}}$ (see Notation 1). We will also use the notation Cl_F^S for $Cl_{F_{i,0}}^{S_{i,0}}$.

Remark 4. Note that if p is odd or F is nonexceptional, $Cl_{F_{i,n}}^{S_{i,n}}$ is also equal to the quotient of the narrow class group of $F_{i,n}$ by the subgroup generated by the classes of finite primes in $S_{i,n}$.

The following result is the main ingredient of the proof of Theorem 1.

Proposition 3 (Tate-Keune-Weibel). *For every $n \in \mathbb{N}$ and every $i \geq 1$ there is an exact sequence of $\Gamma_{i,n}$ -modules as follows*

$$0 \rightarrow Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} \xrightarrow{J_{i,n}} H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes i+1}) \xrightarrow{\text{inv}} \bigoplus_{w|p} \mu_{p^n}^{\otimes i} \xrightarrow{c} \mu_{p^n}^{\otimes i} \rightarrow 0$$

where c is the map of Lemma 6 and inv is the twist by $\mu_{p^n}^{\otimes i}$ of the direct sum of the local invariant maps (for the definition of $J_{i,n}$, see Remark 5 below). Moreover, if p is odd or n is large enough, taking coinvariants by $\Gamma_{i,n}$ gives

$$0 \rightarrow \left(Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}} \xrightarrow{J_{i,n}} H^2(G_S, \mu_{p^n}^{\otimes i+1}) \xrightarrow{\text{inv}} \left(\bigoplus_{w|p} \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}} \xrightarrow{c} (\mu_{p^n}^{\otimes i})_{\Gamma_{i,n}} \rightarrow 0$$

Proof. The first sequence is exact by Proposition 4.1 of [22]. The second follows from the first by taking $\Gamma_{i,n}$ -coinvariants and using Lemma 6, Lemma 7 and $H_1(\Gamma_{i,n}, \mu_{p^n}^{\otimes i}) = 0$ (by Lemma 4). $\square$

Remark 5. We briefly recall the definition of the map $J_{i,n}$ since it will be used later on. To ease notation, set $E_{i,n} = (F_{i,n})_{S_{i,n}}$ (see Notation 1) and denote by $\mathcal{O}_{E_{i,n}}$ the ring of integers of $E_{i,n}$. The first ingredient we need is the isomorphism (see [12], Proposition 8.3.10)

$$f : Cl_{F_{i,n}}^{S_{i,n}}/p^n \xrightarrow{\sim} H^1(G_{S_{i,n}}, \mathcal{O}_E^\times)/p^n$$

which is defined as follows: if $\mathfrak{a}$ is a fractional ideal of $F_{i,n}$ (not divisible by p), then $f([\mathfrak{a}])$ is represented by the continuous 1-cocycle $\sigma \mapsto \sigma(\alpha)/\alpha$ where $\alpha \in E$ is such that $\alpha\mathcal{O}_E = \mathfrak{a}\mathcal{O}_E$. Now there is an exact sequence (see [12], Proposition 8.3.3)

$$0 \rightarrow \mu_{p^n} \rightarrow \mathcal{O}_E^\times \xrightarrow{p^n} \mathcal{O}_E^\times \rightarrow 0$$

Taking $G_{S_{i,n}}$ -cohomology gives an injective map

$$g : H^1(G_{S_{i,n}}, \mathcal{O}_E^\times)/p^n \rightarrow H^2(G_{S_{i,n}}, \mu_{p^n})$$

Set $g \circ f = J'_{i,n}$: then $J_{i,n}$ is just $J'_{i,n}$ twisted by $\mu_{p^n}^{\otimes i}$.

The following two lemmas clarify the connection between $\Omega_{i,n}^{(p)}$ whose precise description is given in Theorem 1.

Lemma 8. *Suppose that p is an odd prime or F is nonexceptional. Then, for each $n \in \mathbb{N}$ and each $i, j \geq 1$, the following diagram is commutative*

$$\begin{array}{ccc} \bigoplus_{w \nmid p^\infty} H^1((k_{i,n})_w, \mathbb{Z}_p(j))[p^n] & \xrightarrow{I_{F_{i,n},j,n}^c} & H^2(G_{S_{i,n}}, \mathbb{Z}_p(j+1))/p^n \\ \uparrow & & \downarrow \\ \bigoplus_{w \nmid p^\infty} H^0((k_{i,n})_w, \mu_{p^n}^{\otimes j}) & \xrightarrow{-T_{F_{i,n},j,n}} & H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes j+1}) \end{array} \quad (7)$$

Here the left-hand and the right vertical maps are the isomorphisms of Lemma 2, the map $I_{i,j,n}^c$ is induced by the snake lemma on the exact sequence (4) and the map $T_{F_{i,n},j,n}$ is the one appearing in the long exact localization sequence

$$0 \rightarrow H^1(G_{S_{i,n}}, \mu_{p^n}^{\otimes j+1}) \rightarrow H^1(F_{i,n}, \mu_{p^n}^{\otimes j+1}) \rightarrow \bigoplus_{w \nmid p\infty} H^0((k_{i,n})_w, \mu_{p^n}^{\otimes j}) \xrightarrow{T_{F_{i,n},j,n}} H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes j+1}) \rightarrow \dots \quad (8)$$

(see [17], Proposition 1).

Proof. It is easy to see, using the results of [19], § 2, that the commutativity of (7) is equivalent to the commutativity of the following diagram

$$\begin{array}{ccc} \bigoplus_{w \nmid p\infty} H^0((k_{i,n})_w, \mathbb{Q}_p/\mathbb{Z}_p(j))[p^n] & \longrightarrow & H^1(G_{S_{i,n}}, \mathbb{Q}_p/\mathbb{Z}_p(j+1))/p^n \\ \uparrow & & \downarrow \\ \bigoplus_{w \nmid p\infty} H^0((k_{i,n})_w, \mu_{p^n}^{\otimes j}) & \xrightarrow{-T_{F_{i,n},j,n}} & H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes j+1}) \end{array} \quad (9)$$

Here the left-hand vertical map is induced by the injection $\mu_{p^n}^{\otimes j} \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(j)$ while the right-hand vertical map is the connecting homomorphisms for the exact sequence

$$0 \rightarrow \mu_{p^n}^{\otimes j+1} \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(j+1) \xrightarrow{p^n} \mathbb{Q}_p/\mathbb{Z}_p(j+1) \rightarrow 0$$

The upper horizontal map is induced by the snake lemma on the exact sequence at the bottom of diagram (3).

Now recall that the exact sequence (8) comes from the Leray spectral sequence

$$H_{\acute{e}t}^s(\mathcal{O}_{F_{i,n}}^{S_{i,n}}, R^t \iota_* \mu_{p^n}^{\otimes j+1}) \Rightarrow H_{\acute{e}t}^{s+t}(F_{i,n}, \mu_{p^n}^{\otimes j+1})$$

where $\iota : \text{Spec}(F_{i,n}) \rightarrow \text{Spec}(\mathcal{O}_{F_{i,n}}^{S_{i,n}})$ is the morphism of schemes induced by $\mathcal{O}_{F_{i,n}}^{S_{i,n}} \hookrightarrow F_{i,n}$. The above spectral sequence can be rewritten as an Hochschild-Serre spectral sequence in Galois cohomology

$$H^s(G_{S_{i,n}}, H^t(H_{S_{i,n}}, \mu_{p^n}^{\otimes j+1})) \Rightarrow H^{s+t}(F_{i,n}, \mu_{p^n}^{\otimes j+1})$$

where $H_{S_{i,n}}$ is the absolute Galois group of $(F_{i,n})_{S_{i,n}}$. Then the map $T_{i,j,n}$ becomes the transgression

$$tg : H^0(G_{S_{i,n}}, H^1(H_{S_{i,n}}, \mu_{p^n}^{\otimes j+1})) \rightarrow H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes j+1})$$

Similarly, the upper horizontal map in (9) becomes the one induced by the snake lemma on the exact sequence

$$0 \rightarrow H^1(G_{S_{i,n}}, \mathbb{Q}_p/\mathbb{Z}_p(j+1))/\text{Div} \rightarrow H^1(F_{i,n}, \mathbb{Q}_p/\mathbb{Z}_p(j+1))/\text{Div} \rightarrow H^0(G_{S_{i,n}}, H^1(H_{i,n}, \mathbb{Q}_p/\mathbb{Z}_p(j+1))) \rightarrow 0$$

(compare with the bottom row of diagram (3)). In other words the commutativity of (9) is equivalent to the commutativity of

$$\begin{array}{ccc} H^0(G_{S_{i,n}}, H^1(H_{S_{i,n}}, \mathbb{Q}_p/\mathbb{Z}_p(j+1))[p^n]) & \longrightarrow & H^1(G_{S_{i,n}}, \mathbb{Q}_p/\mathbb{Z}_p(j+1))/p^n \\ \uparrow & & \downarrow \\ H^0(G_{S_{i,n}}, H^1(H_{S_{i,n}}, \mu_{p^n}^{\otimes j+1})) & \xrightarrow{-tg} & H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes j+1}) \end{array}$$

Using the description of tg given in [12], Proposition (1.6.5), it is not difficult to see that the above diagram is indeed commutative. We give here the explicit computation. Call α the map obtained by composing

the left vertical arrow, the upper horizontal arrow and the right vertical arrow in the above diagram. We have to prove that $\alpha = -tg$. Let $f : H_{S_{i,n}} \rightarrow \mu_{p^n}^{\otimes j+1}$ be a 1-cocycle whose class in $H^1(H_{S_{i,n}}, \mu_{p^n}^{\otimes j+1})$ is fixed by $G_{S_{i,n}}$. Composing f with the natural injection $\mu_{p^n}^{\otimes j+1} \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(j+1)$ we get a 1-cocycle of $H_{S_{i,n}}$ with values in $\mathbb{Q}_p/\mathbb{Z}_p(j+1)$ which we also call f . Clearly the class of f in $H^1(H_{S_{i,n}}, \mathbb{Q}_p/\mathbb{Z}_p(j+1))$ is fixed by $G_{S_{i,n}}$ and annihilated by p^n . Then we can choose a 1-cocycle $f' : G_{F_{i,n}} \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(j+1)$ such that $f'|_{H_{i,n}} = f$ and there exists a 1-cocycle $f'' : G_{S_{i,n}} \rightarrow \mathbb{Q}_p/\mathbb{Z}_p(j+1)$ such that $p^n f' = f'' \pi$ where $\pi : G_{F_{i,n}} \rightarrow G_{S_{i,n}}$ is the natural projection. If $s : G_{S_{i,n}} \rightarrow G_{F_{i,n}}$ is a (continuous) section of π , we have $p^n f' s = f''$. Finally, then, $\alpha([f]) \in H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes j+1})$ is represented by the 2-cocycle

$$(g_1, g_2) \mapsto \partial(f's)(g_1, g_2) = g_1 f' s(g_2) - f' s(g_1 g_2) + f' s(g_1) = -f' s(g_1 g_2) + f'(s(g_1) s(g_2))$$

(the last equality comes from the fact that f' is a cocycle). On the other hand, since any element of $G_{F_{i,n}}$ can be written uniquely as $s(g)h$ for some $g \in G_{S_{i,n}}$ and $h \in H_{S_{i,n}}$, we can define a map $\tilde{f} : G_{F_{i,n}} \rightarrow \mu_{p^n}^{\otimes j+1}$ by $\tilde{f}(s(g)h) = s(g)f(h)$. Then $tg[f] \in H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes j+1})$ is represented by the 2-cocycle (see [12], Proposition (1.6.5) and its proof)

$$\begin{aligned} (g_1, g_2) \mapsto \partial \tilde{f}(s(g_1), s(g_2)) &= s(g_1) \tilde{f}(s(g_2)) - \tilde{f}(s(g_1) s(g_2)) + \tilde{f}(s(g_1)) \\ &= -\tilde{f}(s(g_1) s(g_2)) = -s(g_1 g_2) f' \left(\frac{s(g_1) s(g_2)}{s(g_1 g_2)} \right) \end{aligned}$$

(the last equality comes from the definition of $\tilde{f}$ and the fact that $f'|_{H_{i,n}} = f$). The result follows because f' is a cocycle. $\square$

Remark 6. I wonder if there exists a proof of the preceding lemma which does not make use of explicit calculation. The fact that $-T_{F_{i,n},j,n}$ (and not simply $T_{F_{i,n},j,n}$) is necessary to have a commutative diagram is somehow unexpected.

Lemma 9. *For each $n \in \mathbb{N}$ and each $i \geq 1$, the following diagram is commutative*

$$\begin{array}{ccc} \bigoplus_{w \nmid p^\infty} \mu_{p^n}^{\otimes i} & \xrightarrow{T_{F_{i,n},i,n}} & H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes i+1}) \\ & \searrow \pi_{i,n} \quad \nearrow J_{i,n} & \\ & Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} & \end{array}$$

where, if $\mathfrak{P}_w$ is the ideal representing w in $F_{i,n}$, we set

$$\pi_{i,n}((\zeta_w)_w) = \left[\prod_{w \nmid p^\infty} \mathfrak{P}_w \otimes \zeta_w \right]$$

Proof. The commutativity of the diagram in the statement is equivalent to the commutativity of the following diagram

$$\begin{array}{ccc} \bigoplus_{w \nmid p^\infty} \mathbb{Z}/p^n \mathbb{Z} & \xrightarrow{T_{F_{i,n},0,n}} & H^2(G_{S_{i,n}}, \mu_{p^n}) \\ & \searrow \pi'_{i,n} \quad \nearrow J'_{i,n} & \\ & Cl_{F_{i,n}}^{S_{i,n}} \otimes \mathbb{Z}/p^n \mathbb{Z} & \end{array}$$

where the definition of $\pi'_{i,n}$ is similar to that of $\pi_{i,n}$ and $J'_{i,n}$ is the map of Remark 5. Comparing the Leray spectral sequence and the Hochschild-Serre spectral sequence and using the computation of [17],

III.1.3, we get a commutative diagram

$$\begin{array}{ccc}
\bigoplus_{w \nmid p\infty} \mathbb{Z}/p^n\mathbb{Z} & \xrightarrow{T_{F_{i,n},0,n}} & H^2(G_{S_{i,n}}, \mu_{p^n}) \\
\Sigma \uparrow & & \parallel \\
H^0(G_{S_{i,n}}, H^1(H_{S_{i,n}}, \mu_{p^n})) & \xrightarrow{tg} & H^2(G_{S_{i,n}}, \mu_{p^n})
\end{array}$$

where Σ is the isomorphism defined in the following way. For each $w \nmid p\infty$, let $(F_{i,n})_w$ be a completion of $F_{i,n}$ at w and let $(F_{i,n})_w^{un}$ be the maximal unramified extension of $(F_{i,n})_w$ inside an algebraic closure $\overline{(F_{i,n})_w}$ of $(F_{i,n})_w$. Once an embedding of an algebraic closure of $F_{i,n}$ in $\overline{(F_{i,n})_w}$ has been chosen, we get a restriction map $\rho_w : \text{Gal}(\overline{(F_{i,n})_w}/(F_{i,n})_w^{un}) \rightarrow H_{S_{i,n}}$. Now, given $f \in H^1(H_{S_{i,n}}, \mu_{p^n})^{G_{S_{i,n}}} = \text{Hom}_{G_{S_{i,n}}}(H_{S_{i,n}}, \mu_{p^n})$, we define

$$f \mapsto (f \circ \rho_w)_w \in \bigoplus_{w \nmid p\infty} \text{Hom}(\text{Gal}(\overline{(F_{i,n})_w}/(F_{i,n})_w^{un}), \mu_{p^n}) = \bigoplus_{w \nmid p\infty} H^1((F_{i,n})_w^{un}, \mu_{p^n})$$

The definition of Σ now results via Kummer theory, since

$$H^1((F_{i,n})_w^{un}, \mu_{p^n}) \cong ((F_{i,n})_w^{un})^\times / ((F_{i,n})_w^{un})^{\times p^n} = \langle \pi_w \rangle / \langle \pi_w^{p^n} \rangle = \mathbb{Z}/p^n\mathbb{Z}$$

where $\pi_w \in (F_{i,n})_w$ is any uniformizer. We are then left to prove that $J'_{i,n} \pi'_{i,n} \Sigma = tg$.

Let $f \in H^1(H_{S_{i,n}}, \mu_{p^n})^{G_{S_{i,n}}} = \text{Hom}_{G_{S_{i,n}}}(H_{S_{i,n}}, \mu_{p^n})$ and let $\beta \in F_{S_{i,n}}^\times$ be an element representing f via the Kummer isomorphism $F_{S_{i,n}}^\times / (F_{S_{i,n}}^\times)^{p^n} \cong H^1(H_{S_{i,n}}, \mu_{p^n})$, i.e.

$$f(h) = \frac{h(\beta^{\frac{1}{p^n}})}{\beta^{\frac{1}{p^n}}} \quad \text{for any } h \in H_{S_{i,n}}$$

It is easy to see that $\pi'_{i,n} \Sigma(f) = [\prod_{w \nmid p\infty} \mathfrak{P}_w^{\tilde{w}(\beta)}]$ where $\tilde{w}$ is the extension of the valuation w to $F_{S_{i,n}}$ defined by the chosen embedding $F_{S_{i,n}} \hookrightarrow \overline{(F_{i,n})_w}$. Thanks to the description of $J'_{i,n}$ given in Remark 5, we see that $J'_{i,n} \pi'_{i,n} \Sigma(f) \in H^2(G_{S_{i,n}}, \mu_{p^n})$ is the class represented by the inhomogenous 2-cocycle

$$(\sigma, \tau) \mapsto \sigma \left(\frac{\tau(\beta)}{\beta} \right)^{\frac{1}{p^n}} \cdot \left(\frac{\sigma\tau(\beta)}{\beta} \right)^{-\frac{1}{p^n}} \cdot \left(\frac{\sigma(\beta)}{\beta} \right)^{\frac{1}{p^n}}$$

If $s : G_{S_{i,n}} \rightarrow G_{F_{i,n}}$ is a (continuous) section of the natural projection $G_{F_{i,n}} \rightarrow G_{S_{i,n}}$, this can be rewritten as

$$(\sigma, \tau) \mapsto s(\sigma) \left(\frac{s(\tau)(\beta^{\frac{1}{p^n}})}{\beta^{\frac{1}{p^n}}} \right) \cdot \left(\frac{s(\sigma\tau)(\beta^{-\frac{1}{p^n}})}{\beta^{-\frac{1}{p^n}}} \right) \cdot \left(\frac{s(\sigma)(\beta^{\frac{1}{p^n}})}{\beta^{\frac{1}{p^n}}} \right) = \frac{s(\sigma)s(\tau)(\beta^{\frac{1}{p^n}})}{s(\sigma\tau)(\beta^{\frac{1}{p^n}})}$$

But this is exactly a representative for $tg(f)$ (see the proof of Proposition 1.6.5 in [12]). $\square$

The following lemma is stated in full generality but will be used in the proof of Theorem 1 only for the case when $p = 2$ and F is nonexceptional.

Lemma 10. *Suppose that p is odd or F is nonexceptional. Then for any $i \geq 1$ and any $n \in \mathbb{N}$, $(Cl_{F_{i,n}}^{S_{i,n}})_p = 0$ if and only if the map*

$$(\partial_{F_{i,n},i}^c)|_{H^2(F_{i,n}, \mathbb{Z}_p(i+1))[p^n]} : H^2(F_{i,n}, \mathbb{Z}_p(i+1))[p^n] \longrightarrow \bigoplus_{w \nmid p\infty} H^1((k_{i,n})_w, \mathbb{Z}_p(i))[p^n]$$

is surjective.

Proof. Thanks to Lemma 8,

$$\text{coker} \left((\partial_{F_{i,n},i}^c)_{|H^2(F_{i,n}, \mathbb{Z}_p(i+1))[p^n]} \right) = 0 \iff T_{F_{i,n},i,n} = 0$$

where $T_{F_{i,n},i,n}$ is the map of Lemma 8. Now the statement follows from Lemma 9 (note that the map $\pi_{i,n}$ in the statement of Lemma 9 is surjective). $\square$

Remark 7. An immediate consequence of the preceding lemma is the following: if $\mu_{p^n} \subseteq F$ for some *positive* $n \in \mathbb{N}$ and $(Cl_F^S)_p \neq 0$, then the localization sequence for $K_{2i}(F)_p$ (or equivalently for $H^2(F, \mathbb{Z}_p(i+1))$) does not split for any $i \geq 1$.

We are now ready to state and prove the main result of the paper. As the reader may guess, the case $p = 2$ will require a different approach (and the results obtained are slightly weaker).

Theorem 1. *Let i be a positive integer and let F be a number field.*

- *If p is odd, then for any $n \in \mathbb{N}$ there is an isomorphism*

$$\Omega_{i,n}^{(p)} \cong \left(Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}}$$

In particular, the localization sequence for $K_{2i}(F)_p$ (or equivalently for $H^2(F, \mathbb{Z}_p(i+1))$) splits if and only if for every $n \in \mathbb{N}$ we have $\left(Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}} = 0$.

- *If $p = 2$ and F is nonexceptional, then for $n \in \mathbb{N}$ sufficiently large there is an isomorphism*

$$\Omega_{i,n}^{(2)} \cong \left(Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{2^n}^{\otimes i} \right)_{\Gamma_{i,n}}$$

Moreover the localization sequence for $K_{2i}(F)_2$ (or equivalently for $H^2(F, \mathbb{Z}_2(i+1))$) splits if and only if for every $n \in \mathbb{N}$ we have $\left(Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{2^n}^{\otimes i} \right)_{\Gamma_{i,n}} = 0$.

Proof. For any $n \in \mathbb{N}$, there is a commutative diagram of $\Gamma_{i,n}$ -modules (those on the bottom line have trivial action) with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^2(G_{S_{i,n}}, \mathbb{Z}_p(i+1)) & \longrightarrow & H^2(F_{i,n}, \mathbb{Z}_p(i+1))_p & \xrightarrow{\tilde{\partial}_i^c} & \bigoplus_{w \nmid p} H^1((k_{i,n})_w, \mathbb{Z}_p(i)) \longrightarrow 0 \\ & & \text{\scriptsize } cor_n^{(2)} \downarrow & & \text{\scriptsize } cor_n^{(2)} \downarrow & & \text{\scriptsize } cor_n^{(1)} \downarrow \\ 0 & \longrightarrow & H^2(G_S, \mathbb{Z}_p(i+1)) & \longrightarrow & H^2(F, \mathbb{Z}_p(i+1))_p & \xrightarrow{\partial_i^c} & \bigoplus_{v \nmid p} H^1(k_v, \mathbb{Z}_p(i)) \longrightarrow 0 \end{array}$$

where cor_n is the appropriate (cohomological) corestriction. Following [4], Section 3, we consider a part of the commutative diagram induced by snake lemma, namely

$$\begin{array}{ccc} \bigoplus_{w \nmid p} H^1((k_{i,n})_w, \mathbb{Z}_p(i))[p^n] & \xrightarrow{I_{F_{i,n},i,n}^c} & H^2(G_{S_{i,n}}, \mathbb{Z}_p(i+1))/p^n \\ \downarrow \text{\scriptsize } cor_n^{(1)} & & \downarrow \text{\scriptsize } cor_n^{(2)} \\ \bigoplus_{v \nmid p} H^1(k_v, \mathbb{Z}_p(i))[p^n] & \xrightarrow{I_{F,i,n}^c} & H^2(G_S, \mathbb{Z}_p(i+1))/p^n \end{array}$$

Using Lemma 2, Lemma 3 and Lemma 8 we can rewrite it as follows

$$\begin{array}{ccc}
\bigoplus_{w \nmid p} \mu_{p^n}^{\otimes i} & \xrightarrow{T_{F_{i,n},i,n}} & H^2(G_{S_n}, \mu_{p^n}^{\otimes i+1}) \\
\downarrow \text{cor}_n^{(0)} & & \downarrow \text{cor}_n^{(2)} \\
\bigoplus_{v \nmid p} H^0(k_v, \mu_{p^n}^{\otimes i}) & \xrightarrow{T_{F,i,n}} & H^2(G_S, \mu_{p^n}^{\otimes i+1})
\end{array}$$

Using Lemma 9, we can complete the above diagram to get the following commutative diagram

$$\begin{array}{ccc}
\bigoplus_{w \nmid p} \mu_{p^n}^{\otimes i} & \xrightarrow{T_{F_{i,n},i,n}} & H^2(G_{S_{i,n}}, \mu_{p^n}^{\otimes i+1}) \\
& \searrow \pi_{i,n} & \nearrow J_{i,n} \\
& Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} & \\
\downarrow & & \downarrow \\
\bigoplus_{v \nmid p} H^0(k_v, \mu_{p^n}^{\otimes i}) & \xrightarrow{T_{F,i,n}} & H^2(G_S, \mu_{p^n}^{\otimes i+1})
\end{array}$$

We remark explicitly that $\pi_{i,n}$ is surjective (since elements of the form $[\mathfrak{P}_w] \otimes \zeta_w$, with ζ_w running in $\mu_{p^n}^{\otimes i}$ generate $Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i}$). By definition of $\Omega_{i,n}^{(p)}$ and Lemma 8, we have

$$\text{Im } T_{F,i,n} \cong \text{Im } I_{F,i,n}^c = \Omega_{i,n}^{(p)}$$

Hence, $\eta_{i,n} = \text{cor}_n^{(2)} \circ J_{i,n}$ defines a $\Gamma_{i,n}$ -homomorphism

$$\eta_{i,n} : Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} \rightarrow \Omega_{i,n}^{(p)}$$

which induces a map

$$\eta_{\Gamma_{i,n}} : (Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i})_{\Gamma_{i,n}} \rightarrow \Omega_{i,n}^{(p)}$$

after taking coinvariants.

If p is odd or n is large enough, $\eta_{\Gamma_{i,n}}$ is both injective (this follows from Proposition 3) and surjective (in fact, $\eta_{i,n}$ is surjective because $\pi_{i,n}$ is surjective and $\text{cor}_n^{(1)}$ is surjective by Lemma 5). In particular if p is odd or n is sufficiently large (and F is nonexceptional)

$$(Cl_{F_n}^{S_n} \otimes \mu_{p^n}^{\otimes i})_{\Gamma_n} \cong \Omega_{i,n}^{(p)}$$

and the first statement and part of the second statement of the theorem are proved thanks to Lemma 1. We now focus on the case where $p = 2$ (and of course F is nonexceptional). Suppose first that the localization sequence for $H^2(F, \mathbb{Z}_2(i+1))$ splits, in other words $\Omega_{i,n}^{(2)} = 0$ for any $n \in \mathbb{N}$. Observe that, since $\Gamma_{i,n}$ is a 2-group, Nakayama's lemma implies

$$(Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{2^n}^{\otimes i})_{\Gamma_{i,n}} = 0 \iff (Cl_{F_{i,n}}^{S_{i,n}})_2 = 0 \quad (10)$$

Moreover, if we set $F_{i,\infty} = \cup_{n \in \mathbb{N}} F_{i,n}$, then $F_{i,\infty}/F$ is a $\mathbb{Z}_2$ -extension, since F is nonexceptional. Now, by Lemma 10, we know that $(Cl_F^S)_2 = 0$ (since $\{\pm 1\} \subseteq F$). Moreover we also know that $(Cl_{F_{i,n}}^{S_{i,n}})_2 = 0$ for n large enough, the map $\eta_{\Gamma_{i,n}}$ being an isomorphism for such n . We have to show that $(Cl_{F_{i,n}}^{S_{i,n}})_2 = 0$ for any

$n \in \mathbb{N}$. Let $m \in \mathbb{N}$ be such that $F_{i,m} \neq F$: we can find a prime v above 2 in F which stays inert in $F_{i,m}/F$ because $(Cl_F^S)_2 = 0$ and only primes above 2 can ramify in $\mathbb{Z}_2$ -extensions. In particular there is only one prime above v in $F_{i,\infty}/F$ (since $F_{i,\infty}/F$ is a $\mathbb{Z}_2$ -extension). Now take $n \geq m$ such that $(Cl_{F_{i,n}}^{S_{i,n}})_2 = 0$: $F_{i,n}/F_{i,m}$ has to be disjoint from the 2-split Hilbert class field of F_m since there is a nonsplit prime above 2 in $F_{i,n}/F_{i,m}$. This means that the norm map

$$(Cl_{F_{i,n}}^{S_{i,n}})_2 \longrightarrow (Cl_{F_{i,m}}^{S_{i,m}})_2$$

has to be surjective, which implies $(Cl_{F_{i,m}}^{S_{i,m}})_2 = 0$.

Now suppose that $(Cl_{F_{i,n}}^{S_{i,n}})_2 = 0$ for every $n \in \mathbb{N}$: we prove by induction on n that $\Omega_{i,n}^{(2)} = 0$. Of course we trivially have $\Omega_{i,0}^{(2)} = 0$ (and indeed also $\Omega_{i,1}^{(2)} = 0$ by Lemma 10). Next consider the following commutative diagram

$$\begin{array}{ccc} H^2(F_{i,n}, \mathbb{Z}_2(i+1))[2^n] & \xrightarrow{\partial_{F_{i,n},i}^c} & \bigoplus_{w \nmid p} H^1((k_{i,n})_w, \mathbb{Z}_2(i))[2^n] \\ \downarrow \text{cor}_n^{(2)} & & \downarrow \text{cor}_n^{(1)} \\ H^2(F, \mathbb{Z}_2(i+1))[2^n] & \xrightarrow{\partial_{F,i}^c} & \bigoplus_{v \nmid p} H^1(k_v, \mathbb{Z}_2(i))[2^n] \end{array}$$

A generic element $x = (x_v) \in \bigoplus_{v \nmid 2\infty} H^1(k_v, \mathbb{Z}_2(i))[2^n]$ can be written as a sum $x = y_1 + \dots + y_n$ where, for any $j = 1, \dots, n$, $y_j \in \bigoplus_{v \nmid 2\infty} H^1(k_v, \mathbb{Z}_2(i))[2^n]$ and

$$(y_j)_v = \begin{cases} x_v & \text{if } x_v \in H^1(k_v, \mathbb{Z}_2(i))[2^j] \setminus H^1(k_v, \mathbb{Z}_2(i))[2^{j-1}] \\ 0 & \text{otherwise} \end{cases}$$

By induction, for any $j = 1, \dots, n-1$, there is an element

$$z_j \in H^2(F, \mathbb{Z}_2(i+1))[2^j] \subseteq H^2(F, \mathbb{Z}_2(i+1))[2^n]$$

such that $\partial_{F,i}^c(z_j) = y_j$. Now $(y_n)_v$ is an element of order 2^n in $H^1(k_v, \mathbb{Z}_2(i))[2^n] \cong H^0(k_v, \mu_{2^n}^{\otimes i})$ (by Lemma 2). This means that $H^0(k_v, \mu_{2^n}^{\otimes i}) = \mu_{2^n}^{\otimes i}$ which implies $(k_{i,n})_w = k_v(\mu_{2^n}^{\otimes i}) = k_v$ for any $w|v$ in $F_{i,n}$ (by Lemma 3), i.e. v splits completely in $F_{i,n}/F$. In particular there is an element $w_n \in \bigoplus_{w \nmid 2\infty} H^1((k_{i,n})_w, \mathbb{Z}_2(i))[2^n]$ such that $\text{cor}_n^{(1)}(w_n) = y_n$. Now

$$H^2(F_{i,n}, \mathbb{Z}_2(i+1))[2^n] \xrightarrow{\partial_{F_{i,n},i}^c} \bigoplus_{w \nmid p} H^1((k_{i,n})_w, \mathbb{Z}_2(i))[2^n]$$

is surjective by Lemma 10 and therefore there exists an element $t_n \in H^2(F_{i,n}, \mathbb{Z}_2(i+1))[2^n]$ such that $\partial_{F_{i,n},i}^c(t_n) = w_n$ and clearly $\partial_{F,i}^c(\text{cor}_n^{(2)}(t_n)) = y_n$. This shows that

$$H^2(F, \mathbb{Z}_p(i+1))[2^n] \xrightarrow{\partial_{F,i}^c} \bigoplus_{v \nmid p} H^1(k_v, \mathbb{Z}_p(i))[2^n]$$

is surjective which is equivalent to $\Omega_{i,n}^{(2)} = 0$. □

Remark 8. It is worth noting (and I thank K. Hutchinson for pointing this out to me) that if $\mu_p \subseteq F$, then the splitting criterion of Theorem 1 is independent of i (compare with Remark 7). In fact, in this situation, Γ_n is a (cyclic) p -group and hence by Nakayama's lemma

$$\left(Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}} = 0 \Leftrightarrow Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} = 0 \Leftrightarrow Cl_{F_{i,n}}^{S_{i,n}}/p^n = 0 \Leftrightarrow (Cl_{F_{i,n}}^{S_{i,n}})_p = 0$$

This holds in particular for $p = 2$ and any nonexceptional number field F .

Remark 9. I don't know whether there is an isomorphism $\Omega_{i,n}^{(2)} \cong (Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{2^n}^{\otimes i})_{\Gamma_{i,n}}$ for any $n \in \mathbb{N}$.

The criterion of Theorem 1 is closely related with the triviality of the so-called i -th étale wild kernel of F , which is denoted $WK_{2i}^{\text{ét}}(F)$ and is isomorphic to $\text{div}(K_{2i}(F)_p)$. Recall that, for any odd p and any number field E , if, for any $v \mid p$, E_v denotes a completion of E at v , $WK_{2i}^{\text{ét}}(E)$ is the kernel of the map

$$H^2(G_{E,S}, \mathbb{Z}_p(i+1)) \longrightarrow \bigoplus_{v \mid p} H_{\text{ét}}^2(E_v, \mathbb{Z}_p(i+1))$$

which is induced by, for any $v \mid p$, an embedding of a fixed algebraic closure of E in a fixed algebraic closure of E_v (the kernel of the above map is actually independent of the chosen embedding).

Theorem 2. [Tate, Banaszak-Kolster, Keune, Schneider, Østvær] Suppose that p is an odd prime or F is nonexceptional. For any $i \geq 1$, we have

$$\text{div}(K_{2i}(F)_p) \cong WK_{2i}^{\text{ét}}(F) \cong \varprojlim \left(Cl_{F_{1,n}}^{S_{1,n}} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_n}$$

Proof. The first isomorphism is due to Tate (for $i = 1$ and p odd, see [19]), Banaszak and Kolster (for any i and p odd, see [1], Theorem 3) and Østvær (for any i , $p = 2$ and F non exceptional, see [13], Theorem 9.5). The second isomorphism is due to Keune or Schneider (for p odd, see [8], Theorem 6.6, or [16], §6, Lemma 1) and Østvær (for $p = 2$ and F nonexceptional, see [13], Theorem 6.1). $\square$

Remark 10. As we have already remarked in the introduction of this paper, the condition $\text{div}(K_{2i}(F)_p) = 0$ is certainly necessary for the localization sequence for $K_{2i}(F)_p$ to split since

$$\text{div}(K_{2i}(\mathcal{O}_F)_p) = \text{div} \left(\bigoplus_{v \nmid p\infty} K_{2i-1}(k_v)_p \right) = 0$$

We can check that the criterion of Theorem 1 is indeed consistent with Theorem 2, namely we can give another proof of the fact that, if $\left(Cl_{F_{1,n}}^{S_{1,n}} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}} = 0$ for any $n \in \mathbb{N}$, then $\text{div}(K_{2i}(F)_p) = 0$. This is not difficult thanks to the description given in Theorem 2. For any $i \geq 1$, set

$$F_{i,\infty} = \bigcup_{n \in \mathbb{N}} F_{i,n} \quad \text{and} \quad \Gamma_{i,\infty} = \text{Gal}(F_{i,\infty}/F)$$

Next observe that $F_{i,\infty} \subseteq F_{1,\infty}$ and using the definition of $F_{i,n}$ one easily shows that the restriction gives an isomorphism

$$\Gamma_{1,\infty} = \Gamma_{i,\infty} \times \Delta_{i,\infty}$$

where $\Delta_{i,\infty} = \text{Gal}(F_{1,\infty}/F_{i,\infty}) = \langle \gamma \in \Gamma_{1,\infty} \mid \gamma^i = 1 \rangle$. In particular $\Delta_{i,\infty}$ is finite of order coprime with p (since $\Gamma_{1,n} \cong \mathbb{Z}_p \times \text{Gal}(F(\mu_p)/F)$ and $\mathbb{Z}_p$ has no nontrivial finite subgroups) and acts trivially on $\mathbb{Z}_p(i)$. Now set

$$X'_{i,\infty} = \varprojlim Cl_{F_{i,n}}^{S_{i,n}}$$

and note that

$$(X'_{i,\infty} \otimes \mathbb{Z}_p(i))_{\Gamma_{i,\infty}} = \varprojlim \left(Cl_{F_{i,n}}^{S_{i,n}} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}}$$

We have $(X'_{1,\infty})_{\Delta_{i,\infty}} = X'_{i,\infty}$ since $\Delta_{i,\infty}$ has order coprime with p and therefore

$$\text{div}(K_{2i}(F)_p) \cong (X'_{1,\infty} \otimes \mathbb{Z}_p(i))_{\Gamma_{1,\infty}} = \left((X'_{1,\infty})_{\Delta_{i,\infty}} \otimes \mathbb{Z}_p(i) \right)_{\Gamma_{i,\infty}} = (X'_{i,\infty} \otimes \mathbb{Z}_p(i))_{\Gamma_{i,\infty}}$$

which shows that, if $\left(Cl_{F_{1,n}}^{S_{1,n}} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}} = 0$ for any $n \in \mathbb{N}$, then $\text{div}(K_{2i}(F)_p) = 0$. We will see in the next section (Example 2) that the converse of this statement is not true in general: in other words, for any $i \geq 1$, there exists a prime number p and a number field F such that the localization sequence for $K_{2i}(F)_p$ does not split but $\text{div}(K_{2i}(F)_p) = 0$.

4 Examples.

To begin with we analyze the simplest case, namely $F = \mathbb{Q}$. Let p be an odd prime and let $A_{i,n}$ denote the p -Sylow subgroup of $Cl_{F_{i,n}}^{S_{i,n}}$ where $F_{i,n} = \mathbb{Q}(\mu_{p^n}^{\otimes i})$. Let $K_{1,n}$ be the n -th level of the cyclotomic $\mathbb{Z}_p$ -extension of $\mathbb{Q}$. Set $\Delta_{1,n} = \text{Gal}(F_{1,n}/K_n)$ and for every $j \in \mathbb{Z}$, let $A_{1,n}^{(j)}$ denote the ω^j -component of $A_{1,n}$ where $\omega : \Delta_n \rightarrow \mathbb{Z}_p^\times$ denotes the Teichmüller character. We need the following well known result.

Theorem 3 (Kurihara). *Let p be an odd prime. For any $i \geq 1$,*

$$H^2(G_{\mathbb{Q},S}, \mathbb{Z}_p(i+1)) = 0 \iff A^{(-i)} = 0$$

where A is the p -Sylow of the class group of $\mathbb{Q}(\mu_p)$. In particular, the triviality of $H^2(G_{\mathbb{Q},S}, \mathbb{Z}_p(i+1))$ only depends on the class of i modulo $p-1$.

Proof. See [9], Corollary 1.5. □

Example 1. Let F be the field of rational numbers. Since $\text{Gal}(F_{1,n}/F_{i,n})$ acts trivially on $\mu_{p^n}^{\otimes i}$, for every $n \geq 1$ we have

$$\left(A_{1,n} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{1,n}} = \left((A_{1,n})_{\text{Gal}(F_{1,n}/F_{i,n})} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}} = \left(A_{i,n} \otimes \mu_{p^n}^{\otimes i} \right)_{\Gamma_{i,n}}$$

the last equality coming from the fact that there is only one ramified prime in $F_{1,n}/\mathbb{Q}$ and it is totally ramified (use [20], Proposition 13.22). By Nakayama's lemma

$$\left(\left(A_{1,n} \otimes \mu_{p^n}^{\otimes i} \right)_{\Delta_{1,n}} \right)_{\text{Gal}(K_{1,n}/\mathbb{Q})} = 0 \iff \left(A_{1,n} \otimes \mu_{p^n}^{\otimes i} \right)_{\Delta_{1,n}} = 0$$

Furthermore

$$\left(A_{1,n} \otimes \mu_{p^n}^{\otimes i} \right)_{\Delta_{1,n}} \cong A_{1,n}^{(-i)}$$

Moreover, it is easy to see that for any $n \geq 1$

$$A_{1,n}^{(-i)} = 0 \iff A_{1,1}^{(-i)} = 0$$

Hence the localization sequence for $K_{2i}(\mathbb{Q})_p$ splits if and only if $A_{1,1}^{(-i)}$ is trivial. Therefore, by Theorem 3, the localization sequence for $K_{2i}(\mathbb{Q})_p$ splits if and only if $H^2(G_{\mathbb{Q},S}, \mathbb{Z}_p(i+1)) = 0$ (the latter is equivalent to $K_{2i}(\mathbb{Z})_p = 0$ under the Quillen-Lichtenbaum conjecture). Moreover, it can be easily proved that $WK_{2i}^{\text{ét}}(\mathbb{Q}) \cong H_{\text{ét}}^2(G_{\mathbb{Q},S}, \mathbb{Z}_p(i+1))$: therefore in this case the triviality of $\text{div}(K_{2i}(\mathbb{Q})_p)$ is a necessary and sufficient condition for the localization sequence for $K_{2i}(\mathbb{Q})_p$ to split.

Anyway in general the condition $WK_{2i}^{\text{ét}}(F) = 0$ is weaker than the condition of Theorem 1, as we will show in the next example. First we need the following criterion.

Proposition 4. *Let p be an odd prime and let $F/\mathbb{Q}$ be finite Galois extension such that*

- $\mu_p \subseteq F$;
- $(Cl_F^S)_p \cong \mathbb{Z}/p\mathbb{Z}$;
- *there is only one prime above p in F ;*
- $F(\mu_{p^2})/F$ *is a nontrivial extension where every prime over p is totally split.*

Then, for any $i \geq 1$, $WK_{2i}^{\text{ét}}(F)_p$ is trivial but the localization sequence for $K_{2i}(F)_p$ does not split.

Proof. We are going to use Jaulent's theory of logarithmic classes: for notation and basic results the reader is referred to [5]. For any $F/\mathbb{Q}$ finite and Galois (even not satisfying the hypotheses) we have an exact sequence (see [2], §3)

$$0 \rightarrow \widetilde{Cl}_F(p) \rightarrow \widetilde{Cl}_F \xrightarrow{\varphi} (Cl_F^S)_p \rightarrow \deg_F \mathcal{D}\ell / (\deg_F \mathfrak{p})\mathbb{Z}_p \rightarrow 0 \quad (11)$$

where $\mathfrak{p}$ is any prime of F over p . Moreover

$$\deg_F \mathcal{D}\ell = p[F \cap \widehat{\mathbb{Q}}^c : \mathbb{Q}]\mathbb{Z}_p \quad \text{and} \quad \deg_F \mathfrak{p} = \tilde{f}_{\mathfrak{p}} \cdot \deg p = [F_{\mathfrak{p}} \cap \widehat{\mathbb{Q}}_p^c : \mathbb{Q}_p] \cdot p$$

where $\widehat{\mathbb{Q}}^c$ (resp. $\widehat{\mathbb{Q}}_p^c$) is the cyclotomic $\widehat{\mathbb{Z}}$ -extension of $\mathbb{Q}$ (resp. $\mathbb{Q}_p$). Now we want to compare $[F \cap \widehat{\mathbb{Q}}^c : \mathbb{Q}]$ and $[F_{\mathfrak{p}} \cap \widehat{\mathbb{Q}}_p^c : \mathbb{Q}_p]$. We have $v_p([F \cap \widehat{\mathbb{Q}}^c : \mathbb{Q}]) = 0$ thanks to the first and the fourth hypothesis. The fourth hypothesis also implies that $v_p([F_{\mathfrak{p}} \cap \widehat{\mathbb{Q}}_p^c : \mathbb{Q}_p]) = s \geq 1$. In other words

$$\deg_F \mathcal{D}\ell / (\deg_F \mathfrak{p})\mathbb{Z}_p \cong \mathbb{Z}/p^s\mathbb{Z}$$

with $s \geq 1$. Now the third hypothesis implies $\widetilde{Cl}(p) = 0$ (see [2], Lemma 4): therefore (11) implies that $s = 1$ and $\widetilde{Cl}_F = 0$ because of the second hypothesis. Since $\mu_p \subseteq F$, we can use the isomorphism (see [6], Théorème 3)

$$\mu_p^{\otimes i} \otimes \widetilde{Cl}_F \cong WK_{2i}^{\text{ét}}(F)/pWK_{2i}^{\text{ét}}(F)$$

to deduce that $WK_{2i}^{\text{ét}}(F) = 0$. On the other hand $(Cl_F^S)_p$ is nontrivial, hence Theorem 1 (or already Remark 7) tells us that the localization sequence for $K_{2i}(F)_p$ does not split. $\square$

Example 2. We have to find a number field F and an odd prime p satisfying the hypotheses of Proposition 4. We proceed as follows: we choose an odd prime p and a prime ℓ such that $\ell \equiv 1 \pmod{p}$: this ensures that $\mathbb{Q}(\zeta_{\ell})$ has exactly one subextension of degree p which we call E . Let K be the subextension of degree p of $\mathbb{Q}(\mu_{p^2})$: then EK is an abelian number field whose Galois group is isomorphic to $(\mathbb{Z}/p\mathbb{Z})^2$. Let F' be a subextension of degree p of EK which is different from E and K . Now, if the order of p modulo ℓ is not divisible by p , then E has to be totally split at 3. In particular, EK/F' is totally split at p and $F'/\mathbb{Q}$ has only one prime above p (which is totally ramified). We may then choose $F = F'(\mu_p)$: then the first, the third and the fourth hypotheses of Proposition 4 are satisfied. So we are left to find such a prime ℓ with the additional requirement that $(Cl_F^S)_p$ is cyclic of order p .

Choose $p = 3$ and $\ell = 61$: of course we have $61 \equiv 1 \pmod{3}$ and 3 has order 10 modulo 61. We only have one choice for F' and computations with PARI ([14]) reveal that $F = F'(\mu_3) = \mathbb{Q}(\theta)$ where θ is a root of the polynomial $X^6 - 793X^3 + 226981$. We clearly have only one (totally ramified) prime above 3 in F and furthermore $(Cl_F^S)_3 \cong \mathbb{Z}/3\mathbb{Z}$. Then by Proposition 4, for any $i \geq 1$, we deduce that $WK_{2i}^{\text{ét}}(F)_3 = 0$ but the localization sequence for $K_{2i}(F)_3$ does not split.

Remark 11. (i) It seems reasonable to conjecture that, for any $i \geq 1$ and any rational prime p , there exist infinitely many number fields F such that the localization sequence for $K_{2i}(F)_p$ does not split but $\text{div}(K_{2i}(F)_p) = 0$.

(ii) As we have seen there exist a number field F and a prime p such that (for any $i \geq 1$) $\text{div}(K_{2i}(F)_p) = 0$ but the localization sequence for $K_{2i}(F)_p$ does not split. However, for any number field F , any prime p and any $i \geq 1$, we have $\text{div}(K_{2i}(F)_p) = 0$ if and only if $K_{2i}(F)_p$ is isomorphic to a direct sum of finite cyclic groups. This follows from a theorem of Prüfer (see [7], Theorem 11).

References

- [1] G. BANASZAK, *Generalization of the Moore exact sequence and the wild kernel for higher K-groups*, Compositio Math. 86 no. 3 (1993), 281-305.
- [2] F. DIAZ Y DIAZ AND F. SORIANO-GAFIUK, *Approche algorithmique du groupe des classes logarithmiques*, J. Number Theory 76 (1999), 1-15.

- [3] W.G. DWYER AND E.M. FRIEDLANDER, *Algebraic and étale K-theory*, Trans. Am. Math. Soc. 292 no. 1. (1985), 247-280.
- [4] K. HUTCHINSON, *The 2-Sylow subgroup of the wild kernel of exceptional number fields*, J. Number Theory 87 (2001), 222-238.
- [5] J.-F. JAULENT, *Classes logarithmiques des corps de nombres*, J. Théor. Nombres Bordeaux 6 (1994), 301-325.
- [6] J.-F. JAULENT AND A. MICHEL, *Approche logarithmique des noyaux étales sauvages des corps de nombres*, J. Number Theory 120 no. 1 (2006), 72-91.
- [7] I. KAPLANSKY, *Infinite abelian groups*, University of Michigan Publications n. 2, 1954.
- [8] F. KEUNE, *On the structure of the K_2 of the ring of integers in a number field*, K-Theory 2 no. 5 (1989), 625-645.
- [9] M. KURIHARA, *Some remarks on conjectures about cyclotomic fields and K-groups of $\mathbb{Z}$* , Compositio Math. 81 no. 2 (1992), 223-236.
- [10] J. MILNOR, *Algebraic K-theory and quadratic forms*, Invent. Math. 9 (1969/70), 318-344.
- [11] J. MILNOR, *Introduction to algebraic K-theory*, Princeton University Press and Tokyo University Press, 1971.
- [12] J. NEUKIRCH, A. SCHMIDT, K. WINGBERG, *Cohomology of number fields*, Grundlehren der mathematischen Wissenschaften 323, Springer-Verlag, 2000.
- [13] P. A. ØSTVAER, *Wild kernels at the prime 2*, Acta Arith. 112 no. 2 (2004), 109-130.
- [14] PARI/GP, VERSION 2.3.2, Bordeaux, 2007, <http://pari.math.u-bordeaux.fr/>.
- [15] J. ROGNES AND C. WEIBEL, *Étale descent for two-primary algebraic K-theory of totally imaginary number fields*, K-Theory 16 no. 2 (1999), 101-104.
- [16] P. SCHNEIDER, *Über gewisse Galoiscohomologiegruppen*, Math. Z. 168 no. 2 (1979), 181-205.
- [17] C. SOULÉ, *K-théorie des anneaux d'entiers des corps de nombres et cohomologie étale*, Invent. Math. 55 (1979), 251-295.
- [18] J. TATE, *Symbols in Arithmetic*, in *Actes du Congrès International des Mathématiciens (Nice, 1970)*, Tome 1, Gauthier-Villars 1971, 201-211.
- [19] J. TATE, *Relation between K_2 and Galois cohomology*, Invent. Math. 36 (1976), 257-274.
- [20] L. C. WASHINGTON, *Introduction to cyclotomic fields*, GTM 83, Springer-Verlag 1997.
- [21] C. WEIBEL, *Étale Chern classes at the prime 2*, in *Algebraic K-theory and algebraic topology (Lake Louise, AB, 1991)*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 407, Kluwer Acad. Publ., 1993, 249-286.
- [22] C. WEIBEL, *Higher wild kernels and divisibility in the K-theory of number fields*, J. Pure Appl. Algebra 206 no. 1-2 (2006), 222-244.

LUCA CAPUTO
 School of Mathematical Sciences
 University College Dublin
 Belfield, Dublin 4, Ireland
luca.caputo@ucd.ie